INDEX TO AUTHORS

ABBOT, A. See NOSSAL, ABBOT, and MITCHELL, 263

—. See NOSSAL, ABBOT, MITCHELL, and LUMMUS, 277

ALCOTT, JUDITH. See PARKS, CASAZZA, ALCOTT, and MELNICK, 91

ANDERSON, NORMAN D. See DEMEESTER, ANDERSON, and SHAFFER, 731

ASKONAS, BRIGITTE A. See UNANUE and ASKONAS, 915

AUSTEN, K. FRANK. See ORANGE, VALENTINE, and AUSTEN, 767

AVRAMEAS, STRATIS. See LEDUC, AVRAMEAS, and BOUTEILLE, 109

AYOUB, ELIA M., and MCCARTY, MACLYN. Intraphagocytic β-*N*-acetylglucosaminidase. Properties of the enzyme and its activity on Group A streptococcal carbohydrate in comparison with soil bacillus enzyme, 833

BARRON, BRUCE. See FISCHETTI, BARRON, and ZABRISKIE, 475

BECKER, CARL G., and MURPHY, GEORGE E. The experimental induction of glomerulonephritis like that in man by infection with Group A streptococci, 1

BECKER, ELMER L. See WARD and BECKER, 693

BENACERRAF, BARUJ. See GREEN, PAUL, and BENACERRAF, 43

—. See PAUL, SISKIND, and BENACERRAF, 25

BERTLES, JOHN F. See DÖBLER and BERTLES, 711

BLOTH, BJÖRN, CHESEBRO, BRUCE, and SVEHAG, SVEN-ERIC. Ultrastructural studies of human and rabbit αM-globulins, 749

BLOTH, BJÖRN. See CHESEBRO, BLOTH, and SVEHAG, 399

BOCKHORN, HERMANN. See VOGT, BOCKHORN, KOZIMA, and SASAKI, 867

BORSOS, TIBOR. See HOYER, BORSOS, RAPP, and VANNIER, 589

BOUTEILLE, MICHEL. See LEDUC, AVRAMEAS, and BOUTEILLE, 109

BOYSE, EDWARD A. See OLD, STOCKERT, BOYSE, and KIM, 523

BROOME, J. D. Studies on the mechanism of tumor inhibition by L-asparaginase. Effects of the enzyme on asparagine levels in the blood, normal tissues, and 6C3HED lymphomas. Differences in asparagine formation and utilization in asparaginase-sensitive and -resistant lymphoma cells, 1055

BROWN, BARRY W. See ROWLEY, FITCH, MOSIER, SOLLIDAY, COPPLESON, and BROWN, 983

BROWN, ROBIN. See PAPPENHEIMER and BROWN, 1073

BRUGH, MAX. See SHOTTS, FOSTER, BRUGH, JORDAN, and MCQUEEN, 359

BYERS, VERA S., and SERCARZ, ELI E. The X-Y-Z scheme of immunocyte maturation. IV. The exhaustion of memory cells, 307

CAMPBELL, DAN H. See RITTENBERG and CAMPBELL, 717

CASAZZA, ANNA M. See PARKS, CASAZZA, ALCOTT, and MELNICK, 91

CEROTTINI, JEAN-CHARLES. See MC-

CONAHEY, CEROTTINI, and DIXON, 1003
CHESEBRO, BRUCE, BLOTH, BJÖRN, and SVEHAG, SVEN-ERIC. The ultrastructure of normal and pathological IgM immunoglobulins, 399
—. See BLOTH, CHESEBRO, and SVEHAG, 749
CHRISTIAN, CHARLES L. See PINCUS, HABERKERN, and CHRISTIAN, 819
COCHRANE, CHARLES G., and HAWKINS, DAVID. Studies on circulating immune complexes. III. Factors governing the ability of circulating complexes to localize in blood vessels, 137
—, and MÜLLER-EBERHARD, H. J. The derivation of two distinct anaphylatoxin activities from the third and fifth components of human complement, 371
—. See KNIKER and COCHRANE, 119
COPPLESON, LIONEL W. See ROWLEY, FITCH, MOSIER, SOLLIDAY, COPPLESON, and BROWN, 983
COSTELLO, RICHARD. See DUBOS, SCHAEDLER, and COSTELLO, 783
CURREY, H. L. F., and ZIFF, MORRIS. Suppression of adjuvant disease in the rat by heterologous anti-lymphocyte globulin, 185

DEMEESTER, TOM R., ANDERSON, NORMAN D., and SHAFFER, CHARLES F. The effect of heterologous anti-lymphocyte serum on mouse hematopoietic stem cells, 731
DIXON, FRANK J. See EDGINGTON, GLASSOCK, and DIXON, 555
—. See GLASSOCK, EDGINGTON, WATSON, and DIXON, 573
—. See LAMBERT and DIXON, 507
—. See MCCONAHEY, CEROTTINI, and DIXON, 1003
DÖBLER, JOHANNA, and BERTLES, JOHN F. The physical state of hemoglobin in sickle-cell anemia erythrocytes in vivo, 711
DONALDSON, VIRGINIA H. Mechanisms of activation of C'1 esterase in hereditary angioneurotic edema plasma in vitro. The role of Hageman factor, a clot-promoting agent, 411
DORNER, MARIANNE M. See YOUNT, DORNER, KUNKEL, and KABAT, 633
DUBOS, RENÉ, SCHAEDLER, RUSSELL W., and COSTELLO, RICHARD. Lasting biological effects of early environmental influences. I. Conditioning of adult size by prenatal and postnatal nutrition, 783
—. See SAVAGE, DUBOS, and SCHAEDLER, 67
—. See SERAVALLI and DUBOS, 801
DUNN, PHILIP. See SISKIND, DUNN, and WALKER, 55

EBBESEN, P. Spontaneous amyloidosis in differently grouped and treated DBA/2, BALB/c, and CBA mice and thymus fibrosis in estrogen-treated BALB/c males, 387
EDELMAN, GERALD M. See MARCHALONIS and EDELMAN, 891
EDGINGTON, THOMAS S., GLASSOCK, RICHARD J., and DIXON, FRANK J. Autologous immune complex nephritis induced with renal tubular antigen. I. Identification and isolation of the pathogenetic antigen, 555
—. See GLASSOCK, EDGINGTON, WATSON, and DIXON, 573
ELDERS, M. J., PARHAM, B. A., and HUGHES, E. R. Prevention of wasting disease by pregnancy associated with hypertrophy of the fetal thymus, 649

FISCHETTI, VINCENT A., BARRON, BRUCE, and ZABRISKIE, JOHN B. Studies on streptococcal bacterio-

phages. I. Burst size and intracellular growth of Group A and Group C streptococcal bacteriophages, 475

FISCHETTI, VINCENT A., and ZABRISKIE, JOHN B. Studies on streptococcal bacteriophages. II. Adsorption studies on Group A and Group C streptococcal bacteriophages, 489

FITCH, FRANK W. See ROWLEY, FITCH, MOSIER, SOLLIDAY, COPPLESON, and BROWN, 983

FOSTER, JOHN W. See SHOTTS, FOSTER, BRUGH, JORDAN, and MCQUEEN, 359

FOSTER, ROGER, JR., METCALF, DONALD, and KIRCHMYER, ROSEMARY. Induction of bone marrow colony–stimulating activity by a filterable agent in leukemic and normal mouse serum, 853

FRANK, MICHAEL M., and HUMPHREY, JOHN H. The subunits in rabbit anti-Forssman IgM antibody, 967

FRANZL, ROBERT E., and MCMASTER, PHILIP D. The primary immune response in mice. I. The enhancement and suppression of hemolysin production by a bacterial endotoxin, 1087

—. See MCMASTER and FRANZL, 1109

GELL, PHILIP G. H. See KELUS and GELL, 215

GLASGOW, LOWELL A. See MURPHY and GLASGOW, 1035

GLASSOCK, RICHARD J., EDGINGTON, THOMAS S., WATSON, J. IAN, and DIXON, FRANK J. Autologous immune complex nephritis induced with renal tubular antigen. II. The pathogenetic mechanism, 573

—. See EDGINGTON, GLASSOCK, and DIXON, 555

GOLDSCHNEIDER, IRVING, and MCGREGOR, DOUGLAS D. Migration of lymphocytes and thymocytes in the rat. I. The route of migration from blood to spleen and lymph nodes, 155

GRATE, HELEN E. See HELLMAN and GRATE, 605

GREEN, IRA, PAUL, WILLIAM E., and BENACERRAF, BARUJ. Hapten carrier relationships in the DNP-PLL · foreign albumin complex system: Induction of tolerance and stimulation of cells in vitro, 43

HABERKERN, ROY. See PINCUS, HABERKERN, and CHRISTIAN, 819

HAHN, HELMUT H. See KOZAK, HAHN, LENNARZ, and WOOD, 341

HALBERT, S. P., HOLM, S. E., and THOMPSON, A. Cardiac autoantibodies. I. Immunodiffusion analysis of multiple responses evoked homologously and heterologously, 613

HARRIS, GILMOUR. Antibody production in vitro. I. Single cell studies of the secondary response to sheep erythrocytes, 661

—. Antibody production in vitro. II. Effects of actinomycin D and puromycin on the secondary response to sheep erythrocytes, 675

HAWKINS, DAVID. See COCHRANE and HAWKINS, 137

HELLMAN, SAMUEL, and GRATE, HELEN E. Enhanced erythropoiesis with concomitant diminished granulopoiesis in preirradiated recipient mice. Evidence for a common stem cell, 605

HICKLIN, MARTIN D. See HIRSCH, MURPHY, and HICKLIN, 757

HIRSCH, MARTIN S., MURPHY, FREDERICK A., and HICKLIN, MARTIN D. Immunopathology of lymphocytic choriomeningitis virus infection of newborn mice. Anti-thymocyte serum effects on glomerulonephritis and wasting disease, 757

HOLM, S. E. See HALBERT, HOLM, and THOMPSON, 613

HOYER, LEON W., BORSOS, TIBOR,

RAPP, HERBERT J., and VANNIER, WILTON E. Heterogeneity of rabbit IgM antibody as detected by C'1a fixation, 589
HUGHES, E. R. See ELDERS, PARHAM, and HUGHES, 649
HUMPHREY, JOHN H. See FRANK and HUMPHREY, 967
HURVITZ, DAHLIA. See SCHLESINGER and HURVITZ, 1127

IVÁNYI, J., MALER, M., WUDL, L., and SERCARZ, ELI E. High-dose delay of the immune response. Effect of actinomycin D on continuation of the immune response in vitro, 1149

JERKOFSKY, MARYANN. See RAPP, JERKOFSKY, MELNICK, and LEVY, 77
JORDAN, HELEN E. See SHOTTS, FOSTER, BRUGH, JORDAN, and MCQUEEN, 359

KABAT, ELVIN A. See YOUNT, DORNER, KUNKEL, and KABAT, 633
KANTOR, FRED S. Delayed hypersensitivity. I. Effect of in vitro exposure of cells to antigen upon leukocytic transfer of delayed hypersensitivity, 251
KELUS, ANDREW S., and GELL, PHILIP G. H. Immunological analysis of rabbit anti-antibody systems, 215
KIM, JAE HO. See OLD, STOCKERT, BOYSE, and KIM, 523
KIRCHMYER, ROSEMARY. See FOSTER, METCALF, and KIRCHMYER, 853
KNIKER, WILLIAM T., and COCHRANE, CHARLES G. The localization of circulating immune complexes in experimental serum sickness. The role of vasoactive amines and hydrodynamic forces, 119
KOZAK, MARILYN SUE, HAHN, HELMUT H., LENNARZ, WILLIAM J., and WOOD, W. BARRY, JR. Studies on the pathogenesis of fever. XVI. Purification and further chemical characterization of granulocytic pyrogen, 341
KOZIMA, KENITI. See VOGT, BOCKHORN, KOZIMA, and SASAKI, 867
KUNKEL, HENRY G. See YOUNT, DORNER, KUNKEL, and KABAT, 633

LAMBERT, P. H., and DIXON, FRANK J. Pathogenesis of the glomerulonephritis of NZB/W mice, 507
LEDUC, ELIZABETH H., AVRAMEAS, STRATIS, and BOUTEILLE, MICHEL. Ultrastructural localization of antibody in differentiating plasma cells, 109
LENNARZ, WILLIAM J. See KOZAK, HAHN, LENNARZ, and WOOD, 341
LEVY, BARNET. See RAPP, JERKOFSKY, MELNICK, and LEVY, 77
LILLY, FRANK. The effect of *Histocompatibility-2* type on response to the friend leukemia virus in mice, 465
LUMMUS, Z. See NOSSAL, ABBOT, MITCHELL, and LUMMUS, 277
LUNDSTEDT, CLAUS. See VOLKERT and LUNDSTEDT, 327

MALER, M. See IVÁNYI, MALER, WUDL, and SERCARZ, 1149
MARCHALONIS, JOHN J., and EDELMAN, GERALD M. Phylogenetic origins of antibody structure. III. Antibodies in the primary immune response of the sea lamprey, *Petromyzon marinus*, 891
MCCARTY, MACLYN. See AYOUB and MCCARTY, 833
MCCONAHEY, PATRICIA J., CEROTTINI, JEAN-CHARLES, and DIXON, FRANK J. An approach to the quantitation of immunogenic antigen, 1003

McCULLOCH, E. A. See WU, TILL, SIMINOVITCH, and McCULLOCH, 455
McGREGOR, DOUGLAS D. Bone marrow origin of immunologically competent lymphocytes in the rat, 953
—. See GOLDSCHNEIDER and McGREGOR, 155
McMASTER, PHILIP D., and FRANZL, ROBERT E. The primary immune response in mice. II. Cellular responses of lymphoid tissue accompanying the enhancement or complete suppression of antibody formation by a bacterial endotoxin, 1109
—. See FRANZL and McMASTER, 1087
McQUEEN, JAMES L. See SHOTTS, FOSTER, BRUGH, JORDAN, and McQUEEN, 359
MELNICK, JOSEPH. See PARKS, CASAZZA, ALCOTT, and MELNICK, 91
—. See RAPP, JERKOFSKY,, MELNICK and LEVY 77
METCALF, DONALD. See FOSTER, METCALF, and KIRCHMYER, 853
MIHAESCO, CONSTANTIN, and SELIGMANN, MAXIME. Papain digestion fragments of human IgM globulins, 431
MITCHELL, JUDITH. See NOSSAL, ABBOT, and MITCHELL, 263
—. See NOSSAL, ABBOT, MITCHELL, and LUMMUS, 277
MOEHRING, JOAN M., and MOEHRING, THOMAS J. The response of cultured mammalian cells to diphtheria toxin. II. The resistant cell: enhancement of toxin action by poly-L-ornithine, 541
MOEHRING, THOMAS J. See MOEHRING and MOEHRING, 541
MÖLLER, GÖRAN. Regulation of cellular antibody synthesis. Cellular 7S production and longevity of 7S antigen-sensitive cells in the absence of antibody feedback, 291
MOSIER, DONALD E. See ROWLEY, FITCH, MOSIER, SOLLIDAY, COPPLESON, and BROWN, 983
MÜLLER-EBERHARD, H. J. See COCHRANE and MÜLLER-EBERHARD, 371
MURPHY, BRIAN R., and GLASGOW, LOWELL A. Factors modifying host resistance to viral infection. III. Effect of whole body X-irradiation on experimental encephalomyocarditis virus infection, 1035
MURPHY, FREDERICK A. See HIRSCH, MURPHY, and HICKLIN, 757
MURPHY, GEORGE E. See BECKER and MURPHY, 1

NOSSAL, G. J. V., ABBOT, A., and MITCHELL, JUDITH. Antigens in immunity. XIV. Electron microscopic radioautographic studies of antigen capture in the lymph node medulla, 263
—, ABBOT, A., MITCHELL, JUDITH, and LUMMUS, Z. Antigens in immunity. XV. Ultrastructural features of antigen capture in primary and secondary lymphoid follicles, 277

OLD, LLOYD J., STOCKERT, ELISABETH, BOYSE, EDWARD A., and KIM, JAE HO. Antigenic modulation. Loss of TL antigen from cells exposed to TL antibody. Study of the phenomenon in vitro, 523
ORANGE, ROBERT P., VALENTINE, MARTIN D., and AUSTEN, K. FRANK. Antigen-induced release of slow-reacting substance of anaphylaxis (SRS-A^{rat}) in rats prepared with homologous antibody, 767

PAPPENHEIMER, A. M., JR., and BROWN, ROBIN. Studies on the mode of action of diphtheria toxin. VI. Site of toxin action in living cells, 1073
PARHAM, B. A. See ELDERS, PARHAM, and HUGHES, 649
PARKS, WADE P., CASAZZA, ANNA M., ALCOTT, JUDITH, and MELNICK,

JOSPEH L. Adeno-associated satellite virus interference with the replication of its helper adenovirus, 91

PAUL, WILLIAM E., SISKIND, GREGORY W., and BENACERRAF, BARUJ. Specificity of cellular immune responses. Antigen concentration dependence of stimulation of DNA synthesis in vitro by specifically sensitized cells, as an expression of the binding characteristics of cellular antibody, 25

—. See GREEN, PAUL, and BENACERRAF, 43

PINCUS, THEODORE, HABERKERN, ROY, and CHRISTIAN, CHARLES L. Experimental chronic glomerulitis, 819

RAPP, FRED, JERKOFSKY, MARYANN, MELNICK, JOSEPH L., and LEVY, BARNET. Variation in the oncogenic potential of human adenoviruses carrying a defective SV40 genome (PARA), 77

RAPP, HERBERT J. See HOYER, BORSOS, RAPP, and VANNIER, 589

RITTENBERG, MARVIN B., and CAMPBELL, DAN H. Heterologous carriers in the anamnestic antihapten response, 717

ROWLEY, DONALD A., FITCH, FRANK W., MOSIER, DONALD E., SOLLIDAY, SUSAN, COPPLESON, LIONEL W., and BROWN, BARRY W. The rate of division of antibody-forming cells during the early primary immune response, 983

SASAKI, MASAMICHI. See VOGT, BOCKHORN, KOZIMA, and SASAKI, 867

SAVAGE, DWAYNE C., DUBOS, RENÉ, and SCHAEDLER, RUSSELL W. The gastrointestinal epithelium and its autochthonous bacterial flora, 67

SCHAEDLER, RUSSELL W. See DUBOS, SCHAEDLER, and COSTELLO, 783

. See SAVAGE, DUBOS, and SCHAEDLER, 67

SCHECHTER, ISRAEL. Antigenic competition between polypeptidyl determinants in normal and tolerant rabbits, 237

SCHLESINGER, MICHAEL, and HURVITZ, DAHLIA. Serological analysis of thymus and spleen grafts, 1127

SCHWARZ, M. ROY. The mixed lymphocyte reaction: an in vitro test for tolerance, 879

SELIGMANN, MAXIME. See MIHAESCO and SELIGMANN, 431

SELL, STEWART. Studies on rabbit lymphocytes in vitro. VIII. The relationship between heterozygosity and homozygosity of lymphocyte donor and per cent blast transformation induced by anti-allotype sera, 1139

SERAVALLI, EGILDE, and DUBOS, RENÉ. Lasting biological effects of early environmental influences. II. Lasting depression of weight caused by neonatal contamination, 801

SERCARZ, ELI E. See BYERS and SERCARZ, 307

—. See IVÁNYI, MALER, WUDL, and SERCARZ, 1149

SHAFFER, CHARLES F. See DEMEESTER, ANDERSON, and SHAFFER, 731

SHOTTS, EMMETT B., JR., FOSTER, JOHN W., BRUGH, MAX, JORDAN, HELEN E., and MCQUEEN, JAMES L. An intestinal threadworm as a reservoir and intermediate host for swine influenza virus. A confirmation and amplification of Shope's syndrome, 359

SIMINOVITCH, L. See WU, TILL, SIMINOVITCH, and MCCULLOCH, 455

SISKIND, GREGORY W., DUNN, PHILIP, and WALKER, J. GEOFFREY. Studies

on the control of antibody synthesis. II. Effect of antigen dose and of suppression by passive antibody on the affinity of antibody synthesized, 55

SISKIND, GREGORY W. See PAUL, SISKIND, and BENACERRAF, 25

SOLLIDAY, SUSAN. See ROWLEY, FITCH, MOSIER, SOLLIDAY, COPPLESON, and BROWN, 983

SPITZNAGEL, J. K. See ZEYA and SPITZNAGEL, 927

STOCKERT, ELISABETH. See OLD, STOCKERT, BOYSE, and KIM, 523

SVEHAG, SVEN-ERIC. See BLOTH, CHESEBRO, and SVEHAG, 749

—. See CHESEBRO, BLOTH, and SVEHAG, 399

THOMPSON, A. See HALBERT, HOLM, and THOMPSON, 613

TILL, J. E. See WU, TILL, SIMINOVITCH, and MCCULLOCH, 455

TOP, FRANKLIN H., JR., and WANNAMAKER, LEWIS W. The serum opacity reaction of *Streptococcus pyrogens*. The demonstration of multiple, strain-specific lipoproteinase antigens, 1013

TRENTIN, J. J. See WOLF and TRENTIN, 205

UNANUE, EMIL R., and ASKONAS, BRIGITTE A. Persistence of immunogenicity of antigen after uptake by macrophages, 915

VALENTINE, MARTIN D. See ORANGE, VALENTINE, and AUSTEN, 767

VANNIER, WILTON E. See HOYER, BORSOS, RAPP, and VANNIER, 589

VIROLAINEN, MARTTI. Hematopoietic origin of macrophages as studied by chromosome markers in mice, 943

VOGT, ARNOLD, BOCKHORN, HERMANN, KOZIMA, KENITI, and SASAKI, MASAMICHI. Electron microscopic localization of the nephrotoxic antibody in the glomeruli of the rat after intravenous application of purified nephritogenic antibody-ferritin conjugates, 867

VOLKERT, MOGENS, and LUNDSTEDT, CLAUS. The provocation of latent lymphocytic choriomeningitis virus infections in mice by treatment with anti-lymphocytic serum, 327

WALKER, J. GEOFFREY. See SISKIND, DUNN, and WALKER, 55

WANNAMAKER, LEWIS W. See TOP and WANNAMAKER, 1013

WARD, PETER A., and BECKER, ELMER L. The deactivation of rabbit neutrophils by chemotactic factor and the nature of the activatable esterase, 693

WARNER, NOEL L., and WISTAR, RICHARD, JR. Immunoglobulins in NZB/Bl mice. I. Serum immunoglobulin levels and immunoglobulin class of erythrocyte autoantibody, 169

WATSON, J. IAN. See GLASSOCK, EDGINGTON, WATSON, and DIXON, 573

WISTAR, RICHARD, JR. See WARNER and WISTAR, 169

WOLF, N. S., and TRENTIN, J. J. Hematopoietic colony studies. V. Effect of hematopoietic organ stroma on differentiation of pluripotent stem cells, 205

WOOD, W. BARRY, JR. See KOZAK, HAHN, LENNARZ, and WOOD, 341

WU, A. M., TILL, J. E., SIMINOVITCH, L., and MCCULLOCH, E. A. Cytological evidence for a relationship between normal hematopoietic colony-forming cells and cells of the lymphoid system, 455

WUDL, L. See IVÁNYI, MALER, WUDL, and SERCARZ, 1149

YOUNT, WILLIAM J., DORNER, MARIANNE M., KUNKEL, HENRY G., and KABAT ELVIN, A. Studies on human antibodies. VI. Selective variations in subgroup composition and genetic markers, 633

ZABRISKIE, JOHN B. See FISCHETTI BARRON, and ZABRISKIE, 475

ZABRISKIE, JOHN B. See FISCHETTI, and ZABRISKIE, 489

ZEYA, H. I., and SPITZNAGEL, J. K. Arginine-rich proteins of polymorphonuclear leukocyte lysosomes. Antimicrobial specificity and biochemical heterogenity, 927

ZIFF, MORRIS. See CURREY and ZIFF, 185

INDEX TO SUBJECTS

β-*N*-ACETYLGLUCOSAMINIDASE, intraphagocytic, effects of properties and activity on Group A streptococcal carbohydrate, compared with soil bacillus enzyme, 833

Acid, deoxyribonucleic, synthesis in vitro by sensitized cells, as binding characteristics of cellular antibody, antigen concentration dependence of stimulation of, 25

Actinomycin D, effect on continuation of immune response in vitro, high dose delay, 1149

— and puromycin, effects on secondary response to sheep erythrocytes, 675

Adeno-associated satellite virus interference with replication of helper adenovirus, 91

Adenovirus, helper, and adeno-associated satellite virus interference with replication of, 91

Adenoviruses, human, carrying defective SV40 genome (PARA), variation in oncogenic potential of, 77

Adjuvant disease, suppression by heterologous anti-lymphocyte globulin, 185

Albumin complex system, DNP-PLL·foreign hapten carrier relationships in, induction of tolerance and stimulation of cells in vitro, 43

Allotype, anti-, sera, relationship between heterozygosity and homozygosity of lymphocyte donor and per cent blast transformation induced by, 1139

Amines, vasoactive, and hydrodynamic forces, role of, localization of circulating immune complexes in serum sickness, 119

Amyloidosis, spontaneous, in differently grouped and treated mice and thymus fibrosis in estrogen-treated males, 387

Anamnestic antihapten response, heterologous carriers in, 717

Anaphylatoxin activities derived from third and fifth components of human complement, 371

Anaphylaxis, slow-reacting substance of, antigen-induced release in rats prepared with homologous antibody, 767

Anemia erythrocytes, sickle-cell, in vivo, physical state of hemoglobin in, 711

Angioneurotic edema plasma, hereditary, in vitro, activation of C'1 esterase in, role of Hageman factor, clot-promoting agent, 411

Antiallotype sera, relationship between heterozygosity and homozygosity of lymphocyte donor and per cent blast transformation induced by, 1139

Antibodies, human, selective variations in subgroup composition and genetic markers, 633

— in primary immune response of sea lamprey, *Petromyzon marinus*, phylogenetic origins of antibody structure, 891

Antibody, anti-, systems, immunological analysis of, 215

—, anti-Forssman IgM, subunits in, 967

—, cellular, antigen concentration dependence of stimulation of DNA synthesis in vitro by sensitized cells, as binding characteristics of, 25

— feedback, cellular 7S production

and longevity of 7S antigen-sensitive cells in the absence of, 291
Antibody formation by bacterial endotoxin, cellular response of lymphoid tissue accompanying enhancement or complete suppression of, primary immune response, 1109
—, homologous, antigen-induced release of slow-reacting substance of anaphylaxis in rats prepared with, 767
—, IgM, heterogeneity of, as detected by C'1a fixation, 589
— in differentiating plasma cells, ultrastructural localization of, 109
—, nephrotoxic, in glomeruli after intravenous application of purified nephritogenic antibody-ferritin conjugates, electron microscopic localization of, 867
— production in vitro, effects of actinomycin D and puromycin on secondary response to sheep erythrocytes, 675
— production in vitro, single cell studies of secondary response to sheep erythrocytes, 661
— structure, phylogenetic origins of, antibodies in primary immune response of sea lamprey, *Petromyzon marinus*, 891
— synthesis, cellular 7S production and longevity of 7S antigen-sensitive cells in absence of antibody feedback, 291
— synthesis, effect of antigen dose and of suppression by passive antibody on affinity of, 55
—, TL, loss of TL antigen from cells exposed to, in vitro studies, 523
Antibody-ferritin conjugates, purified nephritogenic, electron microscopic localization of nephrotoxic antibody in glomeruli after intravenous application of, 867
Antibody-forming cells, rate of division during early primary immune response, 983
Antigen after uptake by macrophages, persistence of immunogenicity, 915
— capture in lymph node medulla, electron microscopic radioautographic studies of, 263
— capture in primary and secondary lymphoid follicles, ultrastructural features, 277
— concentration dependence of stimulation of DNA synthesis in vitro by sensitized cells, as binding characteristics of cellular antibody, 25
— dose and suppression by passive antibody, effect on affinity of antibody synthesis, 55
—, immunogenic, quantitation of, 1003
—, in vitro exposure of cells to, effect on leukocytic transfer of delayed hypersensitivity, 251
—, renal tubular, autologous immune complex nephritis induced with, identification and isolation of pathogenetic antigen, 555
—, renal tubular, autologous immune complex nephritis induced with, pathogenetic mechanism, 573
—, TL, loss from cells exposed to TL antibody, in vitro studies, 523
Antigen-induced release of slow-reacting substance of anaphylaxis in rats prepared with homologous antibody, 767
Antigen-sensitive cells, longevity of, and cellular 7S production in absence of antibody feedback, 291
Antigenic competition between polypeptidyl determinants in normal and tolerant rabbits, 237
Antigens, multiple, strain-specific lipoproteinase, serum opacity reaction of *Streptococcus pyogenes*, 1013
Antihapten response, anamnestic, heterologous carriers in, 717
Anti-lymphocyte globulin, heterologous, suppression of adjuvant disease by, 185
— serum, heterologous, effect on hematopoietic stem cells, 731

Anti-lymphocytic serum, provocation of latent lymphocytic choriomeningitis virus infections by treatment with, 327
Antimicrobial specificity and biochemical heterogeneity, arginine-rich proteins of polymorphonuclear leukocyte lysosomes, 927
Anti-thymocyte serum effects on glomerulonephritis and wasting disease, 757
Arginine-rich proteins of polymorphonuclear leukocyte lysosomes, antimicrobial specificity and biochemical heterogeneity, 927
L-Asparaginase, tumor inhibition by, effects of enzyme on asparagine levels in blood, normal tissues, and 6C3HED lymphomas, differences in asparagine formation and utilization in asparaginase-sensitive and -resistant lymphoma cells, 1055
Asparaginase-sensitive and -resistant lymphoma cells, differences in asparagine formation and utilization in, effects of L-asparaginase on asparagine levels in blood, normal tissues, and 6C3HED lymphomas, 1055
Asparagine levels in blood, normal tissues, and 6C3HED lymphomas, effects of L-asparaginase on, differences in asparagine formation and utilization in asparaginase-sensitive and -resistant lymphoma cells, 1055
Autoantibodies, cardiac, immunodiffusion analysis of multiple responses evoked homologously and heterologously, 613
Autoantibody, erythrocyte, immunoglobulin class of, and serum immunoglobulin levels, 169

BACILLUS enzyme, soil, compared with effects of properties and activity of intraphagocytic β-*N*-acetylglucosaminidase on Group A streptococcal carbohydrate, 833
Bacterial endotoxin, cellular response of lymphoid tissue accompanying the enhancement or complete suppression of antibody formation by, primary immune response, 1109
— endotoxin, enhancement and suppression of hemolysin production by, primary immune response, 1087
— flora, autochthonous, and gastrointestinal epithelium, 67
Bacteriophages, streptococcal, Groups A and C, adsorption studies on, 489
—, streptococcal, Groups A and C, burst size and intracellular growth of, 475
Biochemical heterogeneity and antimicrobial specificity, arginine-rich proteins of polymorphonuclear leukocyte lysosomes, 927
Biological effects, lasting, of early environmental influences, conditioning of adult size by prenatal and postnatal nutrition, 783
— effects, lasting, of early environmental influences, lasting depression of weight caused by neonatal contamination, 801
Blast transformation, per cent induced by antiallotype sera, and homozygosity and heterozygosity of lymphocyte donor, relationship between, 1139
Blood, migration route of lymphocytes and thymocytes to spleen and lymph nodes from, 155
—, normal tissues, and 6C3HED lymphomas, effects of L-asparaginase levels in, differences in asparagine formation and utilization in asparaginase-sensitive and -resistant lymphoma cells, 1055
— vessels, factors governing ability of circulating immune complexes to localize in, 137
Bone marrow colony–stimulating activity induced by filterable agent in

leukemic and normal serum, 853
Bone marrow origin of immunologically competent lymphocytes, 953

CARBOHYDRATE, Group A streptococcal, effects of properties and activity of intraphagocytic β-*N*-acetylglucosaminidase on, compared with soil bacillus enzyme, 833
Cardiac autoantibodies, immunodiffusion analysis of multiple responses evoked homologously and heterologously, 613
Cell, intracellular growth and burst size of Group A and Group C streptococcal bacteriophages, 475
—, sickle-, anemia erythrocytes in vivo, physical state of hemoglobin in, 711
—, single, studies of secondary response to sheep erythrocytes, antibody production in vitro, 661
—, stem, evidence for common, enhanced erythropoiesis and diminished granulopoiesis in preirradiated mice, 605
Cells, antibody-forming, rate of division during early primary immune response, 983
—, asparagine-sensitive and -resistant lymphoma, differences in asparagine formation and utilization in, effects of L-asparaginase on asparagine levels in blood, normal tissue, and 6C3HED lymphomas, 1055
—, colony-forming hematopoietic, and cells of the lymphoid system, cytological evidence for a relationship, 455
—, cultured mammalian, response to diphtheria toxin, toxin action enhanced by poly-L-ornithine in resistant cell, 541
—, effect of in vitro exposure to antigen upon leukocytic transfer of delayed hypersensitivity, 251
— exposed to TL antibody, loss of TL antigen from, in vitro studies, 523
Cells, in vitro, induction of tolerance and stimulation of, hapten carrier relationships in the DNP-PLL·foreign albumin complex system, 43
—, living, site of toxin action in, mode of action of diphtheria toxin, 1073
—, memory, exhaustion of, X-Y-Z scheme of immunocyte maturation, 307
—, plasma, differentiating, ultrastructural localization of antibody in, 109
—, sensitized, antigen concentration dependence of stimulation of DNA synthesis in vitro by, as binding characteristics of cellular antibody, 25
—, stem, hematopoietic, effect of heterologous anti-lymphocyte serum on, 731
—, stem, pluripotent, effect of hematopoietic organ stroma on differentiation of, 205
Cellular antibody, antigen concentration dependence of stimulation of DNA synthesis in vitro by sensitized cells, as binding characteristics of, 25
— immune responses, specificity of, 25
— 7S production and longevity of 7S antigen-sensitive cells in absence of antibody feedback, antibody synthesis, 291
— responses of lymphoid tissue accompanying enhancement or complete suppression of antibody formation by bacterial endotoxin, primary immune response, 1109
Chemical characterization and purification of granulocytic pyrogen, 341
Chemotactic factor, deactivation of neutrophils by, and nature of activatable esterase, 693
Choriomeningitis, lymphocytic, virus

infection of newborn mice, immunopathology of, 757
Choriomeningitis virus infections, latent lymphocytic, provocation by treatment with anti-lymphocytic serum, 327
Chromosome markers, hematopoietic origin of macrophages studied by, 943
Clot-promoting agent, role of Hageman factor, activation of C'1 esterase in hereditary angioneurotic edema plasma in vitro, 411
Complement (C'1) esterase, activation in hereditary angioneurotic edema plasma in vitro, role of Hageman factor, clot-promoting agent, 411
— (C'1a) fixation, heterogeneity of IgM antibody as detected by, 589
—, human, anaphylatoxin activities derived from third and fifth components of, 371
Conjugates, antibody-ferritin, purified nephritogenic, electron microscopic localization of nephrotoxic antibody in glomeruli after intravenous application of, 867
Culture, mammalian cells, response to diphtheria toxin, toxin action enhanced by poly-L-ornithine in resistant cell, 541
Cytological evidence for relationship between normal hematopoietic colony-forming cells and cells of lymphoid system, 455

DEOXYRIBONUCLEIC acid synthesis in vitro by sensitized cells, as binding characteristics of cellular antibody, antigen concentration dependence of stimulation of, 25
Digestion, papain, fragments of human IgM globulins, 431
2,4-Dinitrophenyl-poly-L-lysine foreign albumin complex system, hapten carrier relationships, induction of tolerance and stimulation of cells in vitro, 43
Diphtheria toxin, mode of action, site of toxin action in living cells, 1073
Diphtheria toxin, response of cultured mammalian cells to, toxin action enhanced by poly-L-ornithine in resistant cell, 541
Disease, adjuvant, suppression by heterologous anti-lymphocyte globulin, 185
—, wasting and glomerulonephritis, anti-thymocyte serum effects on, 757
—, wasting, prevented by pregnancy associated with hypertrophy of fetal thymus, 649
DNA. *See* Deoxyribonucleic acid.

EDEMA plasma, hereditary angioneurotic, in vitro, activation of C'1 esterase in, role of Hageman factor, clot-promoting agent, 411
Electron microscopic localization of nephrotoxic antibody in glomeruli after intravenous application of purified nephritogenic antibody-ferritin conjugates, 867
— microscopic radioautographic studies of antigen capture in lymph node medulla, 263
Encephalomyocarditis virus infection, effect of whole body X-irradiation on, factors modifying host resistance to viral infection, 1035
Endotoxin, bacterial, cellular response of lymphoid tissue accompanying enhancement or complete suppression of antibody formation by, primary immune response, 1109
—, bacterial, enhancement and suppression of hemolysin production by, primary immune response, 1087
Environmental influences, early, lasting biological effects of, conditioning of adult size by prenatal and postnatal nutrition, 783
— influences, early, lasting biological effects of, lasting depression of

weight caused by neonatal contamination, 801
Enzyme, L-asparaginase, effects on asparagine levels in blood, normal tissues, and 6C3HED lymphomas of mice, differences in asparagine formation and utilization in asparaginase-sensitive and -resistant lymphoma cells, 1055
—, soil bacillus, compared with effects of properties and activity of intraphagocytic β-*N*-acetylglucosaminidase on Group A streptococcal carbohydrate, 833
Epithelium, gastrointestinal, and its autochthonous bacterial flora, 67
Erythrocyte autoantibody, immunoglobulin class of, and serum immunoglobulin levels, 169
Erythrocytes, sheep, effects of actinomycin D and puromycin on secondary response to, 675
—, sheep, single cell studies of secondary response to, antibody production in vitro, 661
—, sickle-cell anemia, in vivo, physical state of hemoglobin in, 711
Erythropoiesis, enhanced, with concomitant diminished granulopoiesis in preirradiated recipient mice, evidence for common stem cell, 605
Esterase, activatable, nature of, and deactivation of neutrophils by chemotactic factor, 693
—, C′1, in hereditary angioneurotic edema plasma in vitro, activation of, role of Hageman factor, clot-promoting agent, 411
Estrogen-treated males, thymus fibrosis in, and spontaneous amyloidosis in differently grouped and treated mice, 387

FERRITIN, antibody-, conjugates, purified nephritogenic, electron microscopic localization of nephrotoxic antibody in glomeruli after intravenous application of, 867
Fetal thymus, prevention of wasting disease by pregnancy associated with hypertrophy of, 649
Fever, pathogenesis of, purification and further chemical characterization of granulocytic pyrogen, 341
Fibrosis, thymus, in estrogen-treated males and spontaneous amyloidosis in differently grouped and treated mice, 387
Flora, bacterial, autochthonous, and gastrointestinal epithelium, 67
Follicles, primary and secondary lymphoid, ultrastructural features of antigen capture in, 277
Forssman IgM antibody, anti-, subunits in, 967
Friend leukemia virus, effect of *Histocompatibility-2* type on response to, 465

GASTROINTESTINAL epithelium and its autochthonous bacterial flora, 67
Genetic markers and subgroup composition in human antibodies, variations, 633
Genome, defective SV40, (PARA), variation in oncogenic potential of human adenoviruses carrying, 77
Globulin, anti-lymphocyte, heterologous, suppression of adjuvant disease by, 185
—, human IgM, papain digestion fragments of, 431
αM-Globulins, human and rabbit, ultrastructural studies of, 749
Glomeruli, electron microscopic localization of nephrotoxic antibody in, after intravenous application of purified nephritogenic antibody-ferritin conjugates, 867
Glomerulitis, chronic, 819
Glomerulonephritis induced by infection with Group A streptococci, 1
—, pathogenesis of, 507
—and wasting disease, anti-thymocyte serum effects on, 757

Glucosaminidase, β-*N*-acetyl-, intraphagocytic, effects of properties and activity on Group A streptococcal carbohydrate, compared with soil bacillus enzyme, 833
Grafts, spleen and thymus, serological analysis of, 1127
Granulocytic pyrogen, purification and further chemical characterization of, 341
Granulopoiesis, diminished, with concomitant erythropoiesis in preirradiated recipient mice, evidence for a common stem cell, 605

HAGEMAN factor, clot-promoting agent, role in activation of C'1 esterase in hereditary edema plasma in vitro, 411
Hapten, anti-, anamnestic, heterologous carriers in, 717
— carrier relationships in DNP-PLL-foreign albumin complex system, induction of tolerance and stimulation of cells in vitro, 43
Hematopoietic colony-forming cells, normal, and cells of lymphoid system, cytological evidence for a relationship, 455
— organ stroma, effect on differentiation of pluripotent stem cells, 205
— origin of macrophages as studied by chromosome markers, 943
— stem cells, effect of heterologous anti-lymphocyte serum on, 731
Hemoglobin in sickle-cell anemia erythrocytes in vivo, physical state of, 711
Hemolysin production by bacterial endotoxin, enhancement and suppression of, primary immune response, 1087
Heterologous anti-lymphocyte globulin, suppression of adjuvant disease by, 185
— anti-lymphocyte serum, effect on hematopoietic stem cells, 731
— carriers in anamnestic antihapten response, 717
Heterozygosity and homozygosity of lymphocyte donor and per cent blast transformation induced by antiallotype sera, relationship between, 1139
Histocompatibility-2 type, effect on response to Friend leukemia virus, 465
Homologous antibody, antigen-induced release of slow-reacting substance of anaphylaxis in rats prepared with, 767
Homozygosity and heterozygosity of lymphocyte donor and per cent blast transformation induced by antiallotype sera, relationship between, 1139
Host resistance to viral infections, effect of whole body X-irradiation on encephalomyocarditis virus infection, 1035
Hydrodynamic forces and vasoactive amines, role of, localization of circulating immune complexes in serum sickness, 119
Hypersensitivity, delayed, effect of in vitro exposure of cells to antigen upon leukocytic transfer of, 251
Hypertrophy of fetal thymus, prevention of wasting disease by pregnancy associated with, 649

IMMUNE complex nephritis, autologous, induced with renal tubular antigen, identification and isolation of pathogenetic antigen, 555
— complex nephritis, autologous, induced with renal tubular antigen, pathogenetic mechanism, 573
— complexes, circulating, factors governing ability to localize in blood vessels, 137
— complexes, circulating, localization in serum sickness, role of vasoactive amines and hydrodynamic forces, 119
— response, early, primary, rate of

division of antibody-forming cells during, 983
Immune response in vitro, high dose delay of, effect of actinomycin D on continuation of, 1149
— response, primary, cellular responses of lymphoid tissue accompanying enhancement or complete suppression of antibody formation by bacterial endotoxin, 1109
— response, primary, enhancement and suppression of hemolysin production by bacterial endotoxin, 1087
— response, primary in sea lamprey, *Petromyzon marinus*, antibodies in, phylogenetic origins of antibody structure, 891
— responses, cellular, specificity of, 25
Immunity, antigens in, electron microscopic radioautographic studies of antigen capture in lymph node medulla, 263
—, antigens in, ultrastructural features of antigen capture in primary and secondary lymphoid follicles, 277
Immunocyte maturation, X-Y-Z scheme of, exhaustion of memory cells, 307
Immunogenic antigen, quantitation of, 1003
Immunogenicity of antigen after uptake by macrophages, persistance of, 915
Immunoglobulin levels, serum, and immunoglobulin class of erythrocyte autoantibody, 169
Immunoglobulins, IgM, ultrastructure of normal and pathological, 399
Immunological analysis of anti-antibody systems, 215
Immunologically competent lymphocytes, bone marrow origin of, 953
Immunopathology of lymphocytic choriomeningitis infection of newborn mice, 757
Infection, virus, encephalomyocarditis, effect of whole body X-irradiation on, factors modifying host resistance to viral infection, 1035
—, virus, lymphocytic, choriomeningitis, of newborn mice, immunopathology of, 757
— with Group A streptococci, induction of glomerulonephritis by, 1
Infections, virus, lymphocytic choriomeningitis, latent, provocation by treatment with anti-lymphocytic serum, 327
Influenza virus, swine, intestinal threadworm as reservoir and intermediate host for, 359
Inhibition, tumor, effect of L-asparaginase on asparagine levels in blood, normal tissues, and 6C3HED lymphomas, differences in asparagine formation and utilization in asparaginase-sensitive and -resistant lymphoma cells, 1055
Injection, intravenous application of purified nephritogenic antibody-ferritin conjugates, electron microscopic localization of nephrotoxic antibody in glomeruli, 867
Intestinal threadworm as reservoir and intermediate host for swine influenza virus, 359
Irradiation, preirradiated recipient mice, enhanced erythropoiesis with concomitant diminished granulopoiesis in, 605

KIDNEY, autologous immune complex nephritis induced with renal tubular antigen, identification and isolation of pathogenetic antigen, 555
—, autologous immune complex nephritis induced with renal tubular antigen, pathogenetic mechanism, 573

LEUKEMIA virus, Friend, effect of *Histocompatibility-2* type on response to, 465

Leukemic and normal serum, induction of bone marrow colony–stimulating activity by filterable agent in, 853
Leukocyte lysosomes, polymorphonuclear, arginine-rich proteins of, antimicrobial specificity and biochemical heterogeneity, 927
Leukocytic transfer of delayed hypersensitivity, effect of in vitro exposure of cells to antigen on, 251
Lipoproteinase antigens, multiple, strain-specific, serum opacity reaction of *Streptococcus pyogenes*, 1013
Lymph node medulla, antigen capture in, electron microscopic radioautographic studies of, 263
— nodes and spleen, migration route of lymphocytes and thymocytes from blood to, 155
Lymphocyte, anti-, globulin, heterologous, suppression of adjuvant disease by, 185
—, anti-, serum, heterologous, effect on hematopoietic stem cells, 731
— donor, heterozygosity and homozygosity of, and per cent blast transformation induced by antiallotype sera, relationship between, 1139
— reaction, mixed, as in vitro test for tolerance, 879
Lymphocytes, immunologically competent, bone marrow origin of, 953
— and thymocytes, migration of, route from blood to spleen and lymph nodes, 155
Lymphocytic choriomeningitis virus infection of newborn mice, immunopathology of, 757
— choriomeningitis virus infections, latent, provocation by treatment with anti-lymphocytic serum, 327
Lymphoid follicles, primary and secondary, ultrastructural features of antigen capture in, 277
— system, cells of, and normal hematopoietic colony-forming cells, cytological evidence for a relationship, 455
Lymphoid tissue, cellular responses of, accompanying enhancement or complete suppression of antibody formation by bacterial endotoxin, primary immune response, 1109
Lymphoma cells, asparaginase-sensitive and -resistant, differences in asparagine formation and utilization in, effects of L-asparaginase on asparagine levels in blood, normal tissues, and 6C3HED lymphomas, 1055
L-Lysine, 2,4-dinitrophenyl-poly-, foreign albumin complex system, hapten carrier relationships in, induction of tolerance and stimulation of cells in vitro, 43
Lysosomes, polymorphonuclear leukocyte, arginine-rich proteins of, antimicrobial specificity and biochemical heterogeneity, 927

MACROPHAGES, hematopoietic origin of, studied by chromosome markers, 943
—, persistance of immunogenicity of antigen after uptake by, 915
Mammalian cells, cultured, response to diphtheria toxin, toxin action enhanced by poly-L-ornithine in resistant cell, 541
Man, adenoviruses carrying defective SV40 genome (PARA), variation in oncogenic potential of, 77
—, antibodies, selective variations in subgroup composition and genetic markers, 633
—, complement, anaphylatoxin activities derived from third and fifth components of, 371
—, glomerulonephritis like that in, induced by infection with Group A streptococci, 1
—, human IgM globulins, papain digestion fragments of, 431
—, human and rabbit αM-globulins, ultrastructural studies of, 749

Marrow, bone, colony–stimulating activity induced by filterable agent in leukemic and normal serum, 853
—, bone, origin of immunologically competent lymphocytes, 953
Medulla, lymph node, antigen capture in, electron microscopic radioautographic studies of, 263
Memory cells, exhaustion of, X-Y-Z scheme of immunocyte maturation, 307
Microbial, anti-, specificity and biochemical heterogeneity, arginine-rich proteins of polymorphonuclear leukocyte lysosomes, 927
Microscopic, electron, localization of nephrotoxic antibody in glomeruli after intravenous application of purified nephritogenic antibody-ferritin conjugates, 867
—, electron, radioautographic studies of antigen capture in lymph node medulla, 263

NEPHRITIS, autologous immune complex, induced with renal tubular antigen, identification and isolation of pathogenetic antigen, 555
—, autologous immune complex, induced with renal tubular antigen, pathogenetic mechanism, 573
—, glomerulo-, induced by infection with Group A streptococci, 1
—, glomerulo-, pathogenesis of, 507
—, glomerulo-, and wasting disease, anti-thymocyte serum effects on, 757
Nephritogenic antibody-ferritin conjugates, purified, electron microscopic localization of nephrotoxic antibody in glomeruli after intravenous application of, 867
Nephrotoxic antibody in glomeruli after intravenous application of purified nephritogenic antibody-ferritin conjugates, electron microscopic localization of, 867
Neutrophils, deactivation by chemotactic factor and nature of activatable esterase, 693
Newborn mice, lymphocytic choriomeningitis virus infection of, immunopathology, 757
Node, lymph, medulla, antigen capture in, electron microscopic radioautographic studies of, 263
Nodes, lymph, route of migration of lymphocytes and thymocytes from blood to spleen and, 155
Nucleic acid, deoxyribo-, synthesis in vitro by sensitized cells, as binding characteristics of cellular antibody, antigen concentration dependence of stimulation of, 25
Nutrition, prenatal and postnatal, conditioning of adult size by, 783

ONCOGENIC potential of human adenoviruses carrying defective SV40 genome (PARA), variation in, 77
Organ stroma, hematopoietic, effect on differentiation of pluripotent stem cells, 205

PAPAIN digestion fragments of human IgM globulins, 431
PARA, defective SV40 genome, variation in oncogenic potential of human adenoviruses carrying, 77
Pathogenesis of fever, purification and further chemical characterization of granulocytic pyrogen, 341
— of glomerulonephritis, 507
Pathogenetic antigen, identification and isolation, autologous immune complex nephritis induced with renal tubular antigen, 555
— mechanism, autologous immune complex nephritis induced with renal tubular antigen, 573
Pathological and normal IgM immunoglobulins, ultrastructure of, 399
Petromyzon marinus, sea lamprey, antibodies in primary response of,

phylogenetic origins of antibody structure, 891
Phagocytic, intraphagocytic β-*N*-acetylglucosaminidase, effects of properties and activity on Group A streptococcal carbohydrate, compared with soil bacillus enzyme, 833
Phylogenetic origins of antibody structure, antibodies in primary immune response of sea lamprey, *Petromyzon marinus*, 891
Plasma cells, differentiating, ultrastructural localization of antibody in, 109
—, hereditary angioneurotic edema, in vitro, activation of C′1 esterase in, role of Hageman factor, clot-promoting agent, 411
Poly-L-ornithine, toxin action enhanced by in resistant cell, response of cultured mammalian cells to diphtheria toxin, 541
Polymorphonuclear leukocyte lysosomes, arginine-rich proteins of, antimicrobial specificity and biochemical heterogeneity, 927
Polypeptidyl determinants in normal and tolerant rabbits, antigenic competition between, 237
Pregnancy associated with hypertrophy of fetal thymus, prevention of wasting disease by, 649
Proteins, arginine-rich, of polymorphonuclear leukocyte lysosomes, antimicrobial specificity and biochemical heterogeneity, 927
Puromycin and actinomycin D, effects on secondary response to sheep erythrocytes, 675
Pyrogen, granulocytic, purification and further chemical characterization of, 341

RADIOAUTOGRAPHIC, electron microscopic, studies of antigen capture in lymph node medulla, 263
Renal. *See* Kidney.
Resistance, asparaginase-resistant and -sensitive lymphoma cells, differences in asparagine formation and utilization in, effects of L-asparaginase on asparagine levels in blood, normal tissues, and 6C3HED lymphomas, 1055
Resistance, host, to viral infections, effect of whole body X-irradiation on encephalomyocarditis virus infection, 1035
Ribonucleic acid, deoxy-, synthesis in vitro by sensitized cells, as binding characteristics of cellular antibody, antigen concentration dependence of stimulation of, 25

SEA lamprey, *Petromyzon marinus*, antibodies in primary immune response of, phylogenetic origins of antibody structure, 891
Sensitive, antigen-, 7S cells, and cellular 7S production, longevity of in absence of antibody feedback, antibody synthesis, 291
Sensitivity, asparaginase-sensitive and -resistant lymphoma cells, differences in asparagine formation and utilization in, effects of L-asparaginase on asparagine levels in blood, normal tissues, and 6C3HED lymphomas, 1055
Sensitized cells, antigen concentration dependence of stimulation of DNA synthesis in vitro by, as binding characteristics of cellular antibody, 25
Sera, antiallotype, relationship between heterozygosity and homozygosity of lymphocyte donor and per cent blast transformation induced by, 1139
Serological analysis of thymus and spleen grafts, 1127
Serum, anti-lymphocyte, heterologous, effect on hematopoietic stem cells, 731
—, anti-lymphocytic, provocation of latent lymphocytic choriomeningitis virus infections by treatment with, 327
—, anti-thymocyte, effects on glomerulonephritis and wasting disease, 757

Serum, immunoglobulin levels and immunoglobulin class of erythrocyte autoantibody, 169
—, leukemic and normal, induction of bone marrow colony–stimulating activity by filterable agent in, 853
— opacity reaction of *Streptococcus pyogenes*, multiple, strain-specific lipoproteinase antigens, 1013
— sickness, localization of circulating immune complexes in, role of vasoactive amines and hydrodynamic forces, 119
Shope's syndrome, confirmation and amplification, intestinal threadworm as reservoir and immediate host for swine influenza virus, 359
Sickle-cell anemia erythrocytes in vivo, physical state of hemoglobin in, 711
Soil bacillus enzyme, compared with effects of properties and activity of intraphagocytic β-*N*-acetylglucosaminidase on Group A streptococcal carbohydrate, 833
Specificity, antimicrobial, and biochemical heterogeneity, arginine-rich proteins of polymorphonuclear leukocyte lysosomes, 927
— of cellular immune responses, 25
—, strain-specific lipoproteinase antigens, multiple, serum opacity reaction of *Streptococcus pyogenes*, 1013
Spleen and lymph nodes, migration route of lymphocytes and thymocytes from blood to, 155
— and thymus grafts, serological analysis of, 1127
Stem cell, common, evidence for, enhanced erythropoiesis with concommitant diminished granulopoiesis in preirradiated mice, 605
— cells, hematopoietic, effect of heterologous anti-lymphocyte serum on, 731
— cells, pluripotent, effect of hematopoietic organ stroma on differentiation of, 205
Stimulation and tolerance of cells in vitro, induction of, hapten carrier relationships in the DNP-PLL·foreign albumin complex system, 43
Streptococcal bacteriophages, Groups A and C, adsorption studies on, 489
— bacteriophages, Groups A and C, burst size and intracellular growth of, 475
— carbohydrate, Group A, effects of properties and activity of intraphagocytic β-*N*-acetylglucosaminidase on, compared with soil bacillus enzyme, 833
Streptococci, Group A, induction of glomerulonephritis by infection with, 1
Streptococcus pyogenes, serum opacity reaction of, multiple, strain-specific lipoproteinase antigens, 1013
Suppression by passive antibody and antigen dose, effect on affinity of antibody synthesis, 55
—, complete, or enhancement of antibody formation by bacterial endotoxin, cellular response of lymphoid tissue accompanying, primary immune response, 1109
— and enhancement of hemolysin production by bacterial endotoxin in primary immune response, 1087
— of adjuvant disease by heterologous anti-lymphocyte globulin, 185
Synthesis, antibody, effect of antigen dose and of suppression by passive antibody on affinity of, 55
—, antibody, in cellular 7S production and longevity of 7S antigen-sensitive cells in absence of antibody feedback, 291
—, DNA, in vitro by sensitized cells, as binding characteristics of cellular antibody, antigen concentration dependence of stimulation of, 25

THREADWORM, intestinal, as reservoir and intermediate host for swine influenza virus, 359
Thymocyte, anti-, serum effects on

glomerulonephritis and wasting disease, 757
Thymocytes and lymphocytes, migration of, route from blood to spleen and lymph nodes, 155
Thymus, fetal, prevention of wasting disease by pregnancy associated with hypertrophy of, 649
— fibrosis in estrogen-treated males and spontaneous amyloidosis in differently grouped and treated mice, 387
— and spleen grafts, serological analysis of, 1127
Tissue, lymphoid, cellular responses of, accompanying enhancement or complete suppression of antibody formation by bacterial endotoxin, primary immune response, 1109
Tissues, normal, blood, and 6C3HED lymphomas of mice, effects of L-asparaginase on asparagine levels in, differences in asparagine formation and utilization in asparaginase-sensitive and -resistant lymphoma cells, 1055
Tolerance, mixed lymphocyte reaction as in vitro test for, 879
— and stimulation of cells in vitro, hapten carrier relationships in the DNP-PLL·foreign albumin complex system, 43
Toxin, diphtheria, mode of action, site of toxin action in living cells, 1073
—, diphtheria, response of cultured mammalian cells to, toxin action enhanced by poly-L-ornithine in resistant cell, 541
Tumor inhibition, effect of L-asparaginase on asparagine levels in blood, normal tissues, and 6C3HED lymphomas, differences in asparagine formation and utilization in asparaginase-sensitive and -resistant lymploma cells, 1055

VIRUS, adeno-associated satellite, interference with replication of helper adenovirus, 91
—, Friend leukemia, effect of *Histocompatibility-2* type on response to, 465
— infection, encephalomyocarditis, effect of whole body X-irradiation on, factors modifying host resistance to viral infection, 1035
— infection, lymphocytic choriomeningitis of newborn mice, immunopathology of, 757
— infections, lymphocytic choriomeningitis, latent, provocation by treatment with anti-lymphocytic serum, 327
—, swine influenza, intestinal threadworm as reservoir and intermediate host for, 359

WASTING disease and glomerulonephritis, anti-thymocyte serum effects on, 757
— disease prevented by pregnancy associated with hypertrophy of fetal thymus, 649
Weight, lasting depression of, caused by neonetal contamination, 801

X-IRRADIATION, whole body, effect on encephalomyocarditis viral infection, factors modifying host resistance to viral infection, 1035